

# 淡江大學九十一學年度碩士班招生考試試題

系別：財務金融學系

科目：統計學

100-1

准帶項目請打「○」否則打「×」	
計算機	字典
○	×

本試題共 6 頁

本試題雙面印製

1. 若  $\hat{y} = 40.768 + 0.1283x$  ( ) 中為標準差  
 (22.14) (0.0305)

若  $y_i^* = y_i / 10$  ,  $x_i^* = x_i / 10$

請列出以下迴歸分析結果，包括標準差。

- (1)  $y$  對  $x^*$  之迴歸分析結果。
- (2)  $y^*$  對  $x^*$  之迴歸分析結果。(10%)

2. 某學者利用新台幣對美元匯率與物價資料進行相對購買力平價關係是否成立之探討，模型為  $\Delta \ln e_t = \alpha_0 + \alpha_1 (\Delta \ln p_t - \Delta \ln p_t^*) + \varepsilon_t$ ， $p_t$  與  $p_t^*$  分別為台灣和美國的躉售物價， $e_t$  為匯率。期間自 1981 年 1 月至 1995 年 6 月。虛無假設為  $H_0: \alpha_0 = 0, \alpha_1 = 1$ 。實證結果如下：

$\alpha_0$	$\alpha_1$
-0.0004	0.8691
(-0.50)	(7.44)

( ) 內為 t 值，請進行假設檢定，在 95% 的顯著水準之下，相對購買力平價關係是否成立？(10%)

3. (1) 設  $y_i = \beta_1 + \beta_2 x_i$ ，樣本數  $T=20$ ，OLS 估計結果如下：

$$\begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \begin{bmatrix} 0.96587 \\ 0.69914 \end{bmatrix}, \text{cov} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \begin{bmatrix} 0.21812 & 0.019195 \\ 0.019195 & 0.048526 \end{bmatrix}, \hat{\sigma}^2 = 2.5193, R^2 = 0.9466$$

請求出  $\beta_2$  之 95% 區間估計值。(5%)

4. 樣本數  $T=63$ ，OLS 估計結果如下：

$$\begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix}, \text{cov} \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 3 & -2 & 1 \\ -2 & 4 & 0 \\ 1 & 0 & 3 \end{bmatrix}$$

在 95% 的顯著水準之下，請進行以下假設檢定： $H_0 = \beta_1 - \beta_2 + \beta_3 = 4$ 。(10%)

5. 某學者進行家計單位消費支出 (C) 每週稅後可支配所得 (Y) 關係的研究，迴歸分析結果如下：

$$\hat{C} = 24.4545 + 0.5091Y \quad ( ) \text{ 內為標準差}$$

(6.4138) (0.0357)

若樣本數  $T=10$ ， $\sum C_i = 1110$ ， $\sum Y_i = 1700$ ， $\sum Y_i C_i = 205500$ ， $\sum Y_i^2 = 32200$ ，

# 淡江大學九十一年度碩士班招生考試試題

系別：財務金融學系

科目：統計學

/00-2

准帶項目請打「○」否則打「×」	
計算機	字典
○	×

本試題共 6 頁

$$\sum C_i^2 = 132100, \sum c_i^2 = 8890, \sum y_i^2 = 33000, \sum y_i c_i = 16800, \bar{C} = 111,$$

$$\bar{Y} = 170, \sum (C_i - \hat{C}_i) = \sum \hat{\varepsilon}_i = 337.2728$$

若未來 A 家庭之週稅後可支配所得  $Y_0 = 100$ ，則該家庭之預測週消費支出  $\hat{C}_0$  為多少？在 5% 顯著水準下，信賴區間為何？(15%)

6. 某學者進行淡水地區房價影響因素之研究，他假設房價 (P) 受坪數 (ST)、臥房數 (RM) 及衛浴數 (BT) 影響，設未受限制之下的迴歸模型為：

$$P = \beta_1 + \beta_2 ST + \beta_3 RM + \beta_4 BT + \varepsilon$$

他懷疑臥房數及衛浴數對房價都不具解釋能力，因此以 F 統計進行假設檢定。已知 URSS (unrestricted residual sum of squares) = 16700, RRSS (restricted residual sum of squares) = 18274, 樣本數  $T = 14$ 。請列出虛無假設與限制下之迴歸模型，然後進行檢定，在 95% 的顯著水準之下，結果如何？(10%)

7. If the salaries of factory workers in a certain industry are approximately normally distributed with a standard deviation of \$500. How large a sample of factory workers would be required if we wish to estimate the population mean salary to within \$60 with a confidence of 99%? (10%)

8. Two methods for packaging a new breakfast cereal produce the same package fill weight. However the second method is slightly faster and will be used unless there is an indication that the variances of the weights produced by the two machines differ. A random sample of 31 packages produced by the second method is analyzed. It is found that the sample standard deviation of weights produced by the first machine is 0.5 ounce while the sample standard deviation of weights produced by the second machine is 0.62 ounce. Find a 90% confidence interval estimate for  $\frac{\sigma_1^2}{\sigma_2^2}$ . (10%)

9. Suppose  $\pi$  is the population proportion, we are interested in testing the null hypothesis  $H_0: \pi \geq 0.25$  with  $\alpha \leq 0.1$  and we wish to detect a difference of 0.025 unit with a minimum power of 0.99. What sample size is required to meet these conditions? (10%)

10. The Superior Research Hospital has publicized the discovery of a new diagnostic test for cancer. If the person has cancer, the test will be *positive* 99% of the time. If the person does not have cancer, the test is *negative* 95% of the time. From past medical

淡江大學九十一學年度碩士班招生考試試題

100-3


系別：財務金融學系

科目：統計學

准帶項目請打「○」否則打「×」	
計算機	字典
○	×

本試題共 6 頁

histories and population census files, the probability that a person has cancer is 0.005. If the test is administered to a randomly selected individual in this population and it is positive, what is the probability that he has cancer? (10%)



Example for  $z = 0.64$   
 $P(0 \leq N(0, 1) \leq 0.64) = 0.2389$

Table 1 Area Under the Standard Normal Distribution

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.0000	0.0040	0.0080	0.0120	0.0160	0.0199	0.0239	0.0279	0.0319	0.0359
0.1	0.0398	0.0438	0.0478	0.0517	0.0557	0.0596	0.0636	0.0675	0.0714	0.0753
0.2	0.0793	0.0832	0.0871	0.0910	0.0948	0.0987	0.1026	0.1064	0.1103	0.1141
0.3	0.1179	0.1217	0.1255	0.1293	0.1331	0.1368	0.1406	0.1443	0.1480	0.1517
0.4	0.1554	0.1591	0.1628	0.1664	0.1700	0.1736	0.1772	0.1808	0.1844	0.1879
0.5	0.1915	0.1950	0.1985	0.2019	0.2054	0.2088	0.2123	0.2157	0.2190	0.2224
0.6	0.2257	0.2291	0.2324	0.2357	0.2389	0.2422	0.2454	0.2486	0.2517	0.2549
0.7	0.2580	0.2611	0.2642	0.2673	0.2704	0.2734	0.2764	0.2794	0.2823	0.2852
0.8	0.2881	0.2910	0.2939	0.2967	0.2995	0.3023	0.3051	0.3079	0.3106	0.3133
0.9	0.3159	0.3186	0.3212	0.3238	0.3264	0.3289	0.3315	0.3340	0.3365	0.3389
1.0	0.3413	0.3438	0.3461	0.3485	0.3508	0.3531	0.3554	0.3577	0.3599	0.3621
1.1	0.3643	0.3665	0.3686	0.3708	0.3729	0.3749	0.3770	0.3790	0.3810	0.3830
1.2	0.3849	0.3869	0.3888	0.3907	0.3925	0.3944	0.3962	0.3980	0.3997	0.4015
1.3	0.4032	0.4049	0.4066	0.4082	0.4099	0.4115	0.4131	0.4147	0.4162	0.4177
1.4	0.4192	0.4207	0.4222	0.4236	0.4251	0.4265	0.4279	0.4292	0.4306	0.4319
1.5	0.4332	0.4345	0.4357	0.4370	0.4382	0.4394	0.4406	0.4418	0.4429	0.4441
1.6	0.4452	0.4463	0.4474	0.4484	0.4495	0.4505	0.4515	0.4525	0.4535	0.4545
1.7	0.4554	0.4564	0.4573	0.4582	0.4591	0.4599	0.4608	0.4616	0.4625	0.4633
1.8	0.4641	0.4649	0.4656	0.4664	0.4671	0.4678	0.4686	0.4693	0.4699	0.4706
1.9	0.4713	0.4719	0.4726	0.4732	0.4738	0.4744	0.4750	0.4756	0.4761	0.4767
2.0	0.4773	0.4778	0.4783	0.4788	0.4793	0.4798	0.4803	0.4808	0.4812	0.4817
2.1	0.4821	0.4826	0.4830	0.4834	0.4838	0.4842	0.4846	0.4850	0.4854	0.4857
2.2	0.4861	0.4864	0.4868	0.4871	0.4875	0.4878	0.4881	0.4884	0.4887	0.4890
2.3	0.4893	0.4896	0.4898	0.4901	0.4904	0.4906	0.4909	0.4911	0.4913	0.4916
2.4	0.4918	0.4920	0.4922	0.4925	0.4927	0.4929	0.4931	0.4932	0.4934	0.4936
2.5	0.4938	0.4940	0.4941	0.4943	0.4945	0.4946	0.4948	0.4949	0.4951	0.4952
2.6	0.4953	0.4955	0.4956	0.4957	0.4959	0.4960	0.4961	0.4962	0.4963	0.4964
2.7	0.4965	0.4966	0.4967	0.4968	0.4969	0.4970	0.4971	0.4972	0.4973	0.4974
2.8	0.4974	0.4975	0.4976	0.4977	0.4977	0.4978	0.4979	0.4979	0.4980	0.4981
2.9	0.4981	0.4982	0.4983	0.4983	0.4984	0.4984	0.4985	0.4985	0.4986	0.4986
3.0	0.4987	0.4987	0.4987	0.4988	0.4988	0.4989	0.4989	0.4989	0.4990	0.4990

Source: This table was generated using the SAS® function PROBNORM

◀ 注意背面尚有試題 ▶

# 淡江大學九十一年度碩士班招生考試試題

系別：財務金融學系

科目：統計學

100-4

准帶項目請打「○」否則打「×」	
計算機	字典
○	×

本試題共 6 頁

TABLE 2  
PERCENTILES OF THE  $\chi^2$  DISTRIBUTION

df	Percent									
	.5	1	2.5	5	10	90	95	97.5	99	99.5
1	.000039	.00016	.00098	.0039	.0158	2.71	3.84	5.02	6.63	7.88
2	.0100	.0201	.0506	.1026	.2107	4.61	5.99	7.38	9.21	10.60
3	.0717	.115	.216	.352	.584	6.25	7.81	9.35	11.34	12.84
4	.207	.297	.484	.711	1.064	7.78	9.49	11.14	13.28	14.86
5	.412	.554	.831	1.15	1.61	9.24	11.07	12.83	15.09	16.75
6	.676	.872	1.24	1.64	2.20	10.64	12.59	14.45	16.81	18.55
7	.989	1.24	1.69	2.17	2.83	12.02	14.07	16.01	18.48	20.28
8	1.34	1.65	2.18	2.73	3.49	13.36	15.51	17.53	20.09	21.96
9	1.73	2.09	2.70	3.33	4.17	14.68	16.92	19.02	21.67	23.59
10	2.16	2.56	3.25	3.94	4.87	15.99	18.31	20.48	23.21	25.19
11	2.60	3.05	3.82	4.57	5.58	17.28	19.68	21.92	24.73	26.76
12	3.07	3.57	4.40	5.23	6.30	18.55	21.03	23.34	26.22	28.30
13	3.57	4.11	5.01	5.89	7.04	19.81	22.36	24.74	27.69	29.82
14	4.07	4.66	5.63	6.57	7.79	21.06	23.68	26.12	29.14	31.32
15	4.60	5.23	6.26	7.26	8.55	22.31	25.00	27.49	30.58	32.80
16	5.14	5.81	6.91	7.96	9.31	23.54	26.30	28.85	32.00	34.27
18	6.26	7.01	8.23	9.39	10.86	25.99	28.87	31.53	34.81	37.16
20	7.43	8.26	9.59	10.85	12.44	28.41	31.41	34.17	37.57	40.00
24	9.89	10.86	12.40	13.85	15.66	33.20	36.42	39.36	42.98	45.56
30	13.79	14.95	16.79	18.49	20.60	40.26	43.77	47.98	50.89	53.67
40	20.71	22.16	24.43	26.51	29.05	51.81	55.76	59.34	63.69	66.77
.60	35.53	37.48	40.48	43.19	46.46	74.40	79.08	83.30	88.38	91.95
120	83.85	86.92	91.58	95.70	100.62	140.23	146.57	152.21	158.95	163.64

Source: Reprinted with permission from W. J. Dixon and F. J. Massey Jr., *Introduction to Statistical Analysis*, 3d ed. (New York: McGraw-Hill, 1969).

TABLE 3  
PERCENTILES OF THE  $t$  DISTRIBUTION

df	P									
	.80	.60	.40	.20	.10	.05	.02	.01		
1	.325	.727	1.376	3.078	6.314	12.706	31.821	63.657		
2	.289	.617	1.061	1.886	2.920	4.303	6.965	9.925		
3	.277	.584	.978	1.638	2.353	3.182	4.541	5.841		
4	.271	.569	.941	1.533	2.132	2.776	3.747	4.604		
5	.267	.559	.920	1.476	2.015	2.571	3.365	4.032		
6	.265	.553	.906	1.440	1.943	2.447	3.143	3.707		
7	.263	.549	.896	1.415	1.895	2.365	2.998	3.499		
8	.262	.546	.889	1.397	1.860	2.306	2.896	3.355		
9	.261	.543	.883	1.383	1.833	2.262	2.821	3.250		
10	.260	.542	.879	1.372	1.812	2.228	2.764	3.169		
11	.260	.540	.876	1.363	1.796	2.201	2.718	3.106		
12	.259	.539	.873	1.356	1.782	2.179	2.681	3.055		
13	.259	.538	.870	1.350	1.771	2.160	2.650	3.012		
14	.258	.537	.868	1.345	1.761	2.145	2.624	2.977		
15	.258	.536	.866	1.341	1.753	2.131	2.602	2.947		
16	.258	.535	.865	1.337	1.746	2.120	2.583	2.921		
17	.257	.534	.863	1.333	1.740	2.110	2.567	2.898		
18	.257	.534	.862	1.330	1.734	2.101	2.552	2.878		
19	.257	.533	.861	1.328	1.729	2.093	2.539	2.861		
20	.257	.533	.860	1.325	1.725	2.086	2.528	2.845		
21	.257	.532	.859	1.323	1.721	2.080	2.518	2.831		
22	.256	.532	.858	1.321	1.717	2.074	2.508	2.819		
23	.256	.532	.858	1.319	1.714	2.069	2.500	2.807		
24	.256	.531	.857	1.318	1.711	2.064	2.492	2.797		
25	.256	.531	.856	1.316	1.708	2.060	2.485	2.787		
26	.256	.531	.856	1.315	1.706	2.056	2.479	2.779		
27	.256	.531	.855	1.314	1.703	2.052	2.473	2.771		
28	.256	.530	.855	1.313	1.701	2.048	2.467	2.763		
29	.256	.530	.854	1.311	1.699	2.045	2.462	2.756		
30	.256	.530	.854	1.310	1.697	2.042	2.457	2.750		
40	.255	.529	.851	1.303	1.684	2.021	2.423	2.704		
60	.254	.527	.848	1.296	1.671	2.000	2.390	2.660		
120	.254	.526	.845	1.289	1.658	1.980	2.358	2.617		
$\infty$	.253	.524	.842	1.282	1.645	1.960	2.326	2.576		

Note: P represents the probability that the  $t$  value will exceed each number in the table in absolute value. This is appropriate for two-tailed tests. For one-tailed tests simply divide each probability in half. For example, .325 in row 1, column 1 tells us that the probability of  $t$  being less than -.325 or greater than .325 is .6.

Source: Obtained from Table III of Fisher and Yates, *Statistical Tables for Biological, Agricultural and Medical Research*, with the permission of the authors and publishers (Edinburgh: Oliver & Boyd, Ltd.).

淡江大學九十一學年度碩士班招生考試試題

100-5

系別：財務金融學系

科目：統計學

准帶項目請打「○」否則打「×」	
計算機	字典
○	×

本試題共 6 頁

Table 4 Right-Tail Critical Values for the F-Distribution

$\alpha$	Upper 1% Points																		
	1	2	3	4	5	6	7	8	9	10	12	15	20	24	30	40	60	120	$\infty$
1	4052.18	4999.50	5403.35	5624.58	5763.65	5858.99	5928.36	5981.07	6022.47	6055.85	6106.32	6157.28	6208.73	6234.63	6260.65	6286.78	6313.03	6339.39	6365.86
2	98.50	99.00	99.17	99.25	99.30	99.33	99.36	99.37	99.39	99.40	99.42	99.43	99.45	99.46	99.47	99.48	99.49	99.49	99.50
3	34.12	30.82	29.46	28.71	28.24	27.91	27.67	27.49	27.35	27.23	27.05	26.87	26.69	26.60	26.50	26.41	26.32	26.22	26.13
4	21.20	18.00	16.69	15.98	15.52	15.21	14.98	14.80	14.66	14.55	14.37	14.20	14.02	13.93	13.84	13.75	13.65	13.56	13.46
5	16.26	13.27	12.06	11.39	10.97	10.67	10.46	10.29	10.16	10.05	9.98	9.92	9.85	9.77	9.72	9.68	9.65	9.62	9.59
6	13.75	10.92	9.78	9.15	8.75	8.47	8.26	8.10	7.98	7.87	7.77	7.72	7.66	7.61	7.57	7.54	7.51	7.48	7.45
7	12.25	9.55	8.45	7.85	7.46	7.19	6.99	6.84	6.72	6.62	6.52	6.47	6.41	6.37	6.34	6.31	6.28	6.25	6.22
8	11.26	8.65	7.59	7.01	6.63	6.37	6.18	6.03	5.91	5.81	5.72	5.67	5.61	5.57	5.54	5.51	5.48	5.45	5.42
9	10.56	8.02	6.99	6.42	6.06	5.80	5.61	5.47	5.35	5.26	5.11	5.06	5.00	4.96	4.93	4.90	4.87	4.84	4.81
10	10.04	7.56	6.55	5.99	5.64	5.39	5.20	5.06	4.94	4.85	4.71	4.66	4.60	4.56	4.53	4.50	4.47	4.44	4.41
11	9.53	7.21	6.22	5.67	5.32	5.07	4.89	4.74	4.63	4.54	4.40	4.35	4.29	4.25	4.22	4.19	4.16	4.13	4.10
12	9.07	6.70	5.74	5.21	4.86	4.62	4.44	4.30	4.19	4.10	3.96	3.92	3.86	3.82	3.79	3.76	3.73	3.70	3.67
13	8.66	6.36	5.42	4.89	4.54	4.30	4.12	4.00	3.89	3.80	3.66	3.62	3.56	3.52	3.49	3.46	3.43	3.40	3.37
14	8.29	6.01	5.09	4.57	4.22	3.98	3.80	3.69	3.60	3.51	3.37	3.33	3.27	3.23	3.20	3.17	3.14	3.11	3.08
15	8.02	5.78	4.87	4.35	4.00	3.76	3.58	3.47	3.38	3.29	3.15	3.11	3.05	3.01	2.98	2.95	2.92	2.89	2.86
16	7.88	5.66	4.76	4.24	3.89	3.65	3.47	3.36	3.27	3.18	3.04	3.00	2.94	2.90	2.87	2.84	2.81	2.78	2.75
17	7.77	5.57	4.68	4.16	3.81	3.57	3.39	3.28	3.19	3.10	2.96	2.92	2.86	2.82	2.79	2.76	2.73	2.70	2.67
18	7.68	5.49	4.60	4.08	3.73	3.49	3.31	3.20	3.11	3.02	2.88	2.84	2.78	2.74	2.71	2.68	2.65	2.62	2.59
19	7.60	5.42	4.54	4.02	3.67	3.43	3.25	3.14	3.05	2.96	2.82	2.78	2.72	2.68	2.65	2.62	2.59	2.56	2.53
20	7.53	5.36	4.48	3.96	3.61	3.37	3.19	3.08	2.99	2.90	2.76	2.72	2.66	2.62	2.59	2.56	2.53	2.50	2.47
21	7.47	5.30	4.42	3.90	3.55	3.31	3.13	3.02	2.93	2.84	2.70	2.66	2.60	2.56	2.53	2.50	2.47	2.44	2.41
22	7.42	5.25	4.37	3.85	3.50	3.26	3.08	2.97	2.88	2.79	2.65	2.61	2.55	2.51	2.48	2.45	2.42	2.39	2.36
23	7.37	5.20	4.32	3.80	3.45	3.21	3.03	2.92	2.83	2.74	2.60	2.56	2.50	2.46	2.43	2.40	2.37	2.34	2.31
24	7.32	5.15	4.27	3.75	3.40	3.16	2.98	2.87	2.78	2.69	2.55	2.51	2.45	2.41	2.38	2.35	2.32	2.29	2.26
25	7.27	5.10	4.22	3.70	3.35	3.11	2.93	2.82	2.73	2.64	2.50	2.46	2.40	2.36	2.33	2.30	2.27	2.24	2.21
26	7.22	5.05	4.17	3.65	3.30	3.06	2.88	2.77	2.68	2.59	2.45	2.41	2.35	2.31	2.28	2.25	2.22	2.19	2.16
27	7.17	5.00	4.12	3.60	3.25	3.01	2.83	2.72	2.63	2.54	2.40	2.36	2.30	2.26	2.23	2.20	2.17	2.14	2.11
28	7.12	4.95	4.07	3.55	3.20	2.96	2.78	2.67	2.58	2.49	2.35	2.31	2.25	2.21	2.18	2.15	2.12	2.09	2.06
29	7.07	4.90	4.02	3.50	3.15	2.91	2.73	2.62	2.53	2.44	2.30	2.26	2.20	2.16	2.13	2.10	2.07	2.04	2.01
30	7.02	4.85	3.97	3.45	3.10	2.86	2.68	2.57	2.48	2.39	2.25	2.21	2.15	2.11	2.08	2.05	2.02	1.99	1.96
40	7.08	4.98	4.13	3.65	3.30	3.06	2.88	2.77	2.68	2.59	2.45	2.41	2.35	2.31	2.28	2.25	2.22	2.19	2.16
60	7.08	4.98	4.13	3.65	3.30	3.06	2.88	2.77	2.68	2.59	2.45	2.41	2.35	2.31	2.28	2.25	2.22	2.19	2.16
120	6.85	4.79	3.95	3.48	3.13	2.89	2.71	2.60	2.51	2.42	2.28	2.24	2.18	2.14	2.11	2.08	2.05	2.02	1.99
$\infty$	6.63	4.61	3.78	3.32	2.97	2.73	2.55	2.44	2.35	2.26	2.12	2.08	2.02	1.98	1.95	1.92	1.89	1.86	1.83

Source: This table was generated using the SAS® function FINV.  $\nu_1$  = numerator degrees of freedom;  $\nu_2$  = denominator degrees of freedom

◀ 注意背面尚有試題 ▶

5

# 淡江大學九十一年度碩士班招生考試試題


100-6

系別：財務金融學系

科目：統計學

准帶項目請打「○」否則打「×」	
計算機	字典
○	×

本試題共 頁



**Table 5 Right-Tail Critical Values for the F-Distribution**

$\nu_1 \backslash \nu_2$	1	2	3	4	5	6	7	8	9	10	12	15	20	24	30	40	60	120	$\infty$
1	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54	241.88	243.91	245.95	248.01	249.05	250.10	251.14	252.20	253.25	254.31
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38	19.40	19.41	19.43	19.45	19.45	19.46	19.47	19.48	19.49	19.50
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.70	8.66	8.64	8.62	8.59	8.57	8.55	8.53
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.86	5.80	5.77	5.75	5.72	5.69	5.66	5.63
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.68	4.62	4.56	4.53	4.50	4.46	4.43	4.40	4.37
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.00	3.94	3.87	3.84	3.81	3.77	3.74	3.70	3.67
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.57	3.51	3.44	3.41	3.38	3.34	3.30	3.27	3.23
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.28	3.22	3.15	3.12	3.08	3.04	3.01	2.97	2.93
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.07	3.01	2.94	2.90	2.86	2.83	2.79	2.75	2.71
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.91	2.85	2.77	2.74	2.70	2.66	2.62	2.58	2.54
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.79	2.72	2.65	2.61	2.57	2.53	2.49	2.45	2.40
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.69	2.62	2.54	2.51	2.47	2.43	2.38	2.34	2.30
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.60	2.53	2.46	2.42	2.38	2.34	2.30	2.25	2.21
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.53	2.46	2.39	2.35	2.31	2.27	2.22	2.18	2.13
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.48	2.40	2.33	2.29	2.25	2.20	2.16	2.11	2.07
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	2.49	2.42	2.35	2.28	2.24	2.19	2.15	2.11	2.06	2.01
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	2.45	2.38	2.31	2.23	2.20	2.15	2.10	2.06	2.01	1.96
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	2.41	2.34	2.27	2.19	2.15	2.11	2.06	2.02	1.97	1.92
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	2.38	2.31	2.23	2.16	2.11	2.07	2.03	1.98	1.93	1.88
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	2.35	2.28	2.20	2.12	2.08	2.04	1.99	1.95	1.90	1.84
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37	2.32	2.25	2.18	2.10	2.05	2.01	1.96	1.92	1.87	1.81
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	2.30	2.23	2.15	2.07	2.03	1.98	1.94	1.89	1.84	1.78
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	2.27	2.20	2.13	2.05	2.01	1.96	1.91	1.86	1.81	1.76
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.18	2.11	2.03	1.98	1.94	1.89	1.84	1.79	1.73
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	2.24	2.16	2.09	2.01	1.96	1.92	1.87	1.82	1.77	1.71
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	2.22	2.15	2.07	1.99	1.95	1.90	1.85	1.80	1.75	1.69
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25	2.20	2.13	2.06	1.97	1.93	1.88	1.84	1.79	1.73	1.67
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	2.19	2.12	2.04	1.96	1.91	1.87	1.82	1.77	1.71	1.65
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22	2.18	2.10	2.03	1.94	1.90	1.85	1.81	1.75	1.70	1.64
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.09	2.01	1.93	1.89	1.84	1.79	1.74	1.68	1.62
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.00	1.92	1.84	1.79	1.74	1.69	1.64	1.58	1.51
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	1.99	1.92	1.84	1.75	1.70	1.65	1.59	1.53	1.47	1.39
120	3.92	3.07	2.68	2.45	2.29	2.18	2.09	2.02	1.96	1.91	1.83	1.75	1.66	1.61	1.55	1.50	1.43	1.35	1.25
$\infty$	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88	1.83	1.75	1.67	1.57	1.52	1.46	1.39	1.32	1.22	1.10

Source: This table was generated using the SAS® function FINV.  $\nu_1$  = numerator degrees of freedom;  $\nu_2$  = denominator degrees of freedom